

WELL CHILD EXAM - EARLY CHILDHOOD 2 YEARS

(Meets EPSDT Guidelines)

童年早期:2歳					
	兒童姓名	兒童由誰帶來	出生日期		
由家長填寫兒童的情況	過敏症	目前用藥			
	自上次體檢後出現的病症/事故/問題/擔憂	今天我有	与一個不明白的問題,關於:		
	是 否 我的孩子飲食多樣化。 我的孩子能順利就寢,並且睡眠很好。 我的孩子能關球。	是 否 我的孩子能搭積木。 我的孩子能講有兩三 我的孩子對大小便訓	E個詞的句子。		
WEIGHT K	G./OZ. PERCENTILE HEIGHT CM/IN. PERCENTILE HEAD CIR. PERC				
R	eview of systems Review of family history				
Screening: N Hearing Vision Development: Circle area of concern Adaptive/Cognitive Language/Communication Gross Motor Social/Emotional Fine Motor Behavior Mental Health Physical: General appearance Skin Lungs Head Cardiovascular/Pulses Eyes Cover/Uncover Ears Genitalia Nose Oropharynx/Teeth Neck Neurologic Nodes Describe abnormal findings and comments.		Review Immunization Fluoride Supplements Hct/Hgb Health Education: (Check Nutrition Toil Development Passive Smoking Child Care Other:	Development Car Seat or Booster Seat TV Habits Passive Smoking Discipline/Limits Teeth Care Child Care Other:		
		IMMUNIZATIONS GIVEN			
		REFERRALS			
NEXT VISIT: 3 YEARS OF AGE			HEALTH PROVIDER NAME		
	PROVIDER SIGNATURE	HEALTH PROVIDER ADDRE			
มอศ ธ 13-68	4A CH (REV. 08/2004)		SEE DICTATION		

DISTRIBUTION: White - Physician Yellow - Parent/Guardian

Guidance to Physicians and Nurse Practitioners for Early Childhood (2 years)

The following highlight EPSDT screens where practitioners often have questions. They are not comprehensive guidelines.

Fluoride Screen

Fluoride supplements should be considered for all children drinking fluoride deficient (<0.6 ppm F) water. Before supplements are prescribed, it is essential to know the fluoride concentration of the patient's drinking water. Once the fluoride level of the water supply has been evaluated, either through contacting the public health officials or water analysis, as well as evaluating other sources of fluoride, the daily dosage schedule can be recommended. Pediatric Dentistry: Reference Manual 1999--00.(21)5.

Lead Screen

Screen children for these risk factors:

- Live in or frequently visits day care center, preschool, baby sitter's home or other structure built before 1950 that is dilapidated or being renovated.
- Come in contact with other children with known lead toxicity (i.e., blood lead 15 ug/dl).
- Live near a lead processing plant or with parents or household members who work in a lead-related occupation (e.g., battery recycling plant).

Pneumococcal Screen

Screen children for these risk factors:

 Age 2 and over with chronic illnesses specifically associated with pneumococcal disease or its complications; anatomic or functional aspenia; sickle cell disease; nephritic syndrome or chronic renal failure; cerebrospinal fluid leaks; or conditions associated with immunosuppression.

Hepatitis B Vaccine

• For children and adolescents not vaccinated against hepatitis B in infancy, begin the hepatitis B vaccine series during any childhood visit. Give the second dose at least one month after the first dose and give the third dose at least four months after the first dose and at least two months after the second dose.

Developmental Milestones

Always ask about and follow-up on parent concerns about development or behavior. You may use the following screening list, or use the Ages and Stages Questionnaire, the Denver II, the ELMS2 (a language screen), or the MacArthur Communication Development Inventory.

Yes	No				
		Kicks ball forward.			
		Removes article of clothing (not hat).			
		Combines 2 words.			
		Uses own name to refer to self.			
		Vocabulary of more than 50 words.			
		Strangers understand half child's speech.			
		Understands a two-step verbal command ("Pick up the toy; put it away") without gestures.			
		Points to 6 named body parts (nose, eyes, ears, mouth, hands, feet, tummy, hair).			
		Stacks 4 blocks.			
Persistent rocking, hand flapping, head banging, or toe walking.					
No spontaneous two word (not echolalic) phrases.					
• ,	Any loss of any language or social skills.				

Instructions for developmental milestones: At least 90% of children should achieve <u>the underlined milestones</u> by this age. If you have checked "no" on **any two items**, *even one* of the underlined items, or any of the **boxed items** (abnormal behavior at this age), refer the child for a formal developmental assessment.

Notes: Immunization schedules are from the Advisory Committee on Immunization Practice of the U.S. Centers for Disease Control and Prevention. Parents and providers may call Healthy Mothers, Healthy Babies (1-800-322-2588) with questions or concerns on childhood development.



健康兒童體檢 - 童年早期: 2歳 WELL CHILD EXAM - EARLY CHILDHOOD 2 YEARS

(符合EPSDT早期定期檢查、診斷和治療指南標準)

日期		

童年早期:2歳							
	兒童姓名		兒童由	誰帶來			出生日期
r-2	過敏症				目前用藥		
的情汤	自上次體檢後出現的病症/事故/問題/擔憂		今天我有一個不明白的問題,關於:		問題,關於:		
寫兒童							
由家長填寫兒童的情況	我的孩子能順利就寢,並且睡眠很好。 我的孩子能講有兩三個詞的句子。 我的孩子能踢球。 我的孩子對大小便訓練感興趣。				0		
體重 千克/	盎司 百分比	身高 厘米/英寸 百分比 遅	圍 百分比	飲	食		
	統審核	家族健康史審核		排	排洩		
				睡眠 審核免疫記錄 鉛中毒現象			
檢查:		E常 不正常		氟化物藥品 氟化物鈾劑			
聽力檢				保保	□ Hct/Hgb □ TB(肺結核) □ 牙科轉介 保健教育:(勾劃所有已討論過/已發過相關資料的項目)		
	發育:請勾劃出您感到擔心的方面				管養 大小便訓練 安全		
適應/認 大的肌	8知 肉運動 社交/	語言/ 交流溝通 情緒 細致的肌	肉運動		一 發育 一 兒童車座椅或加固兒童車座 一 看電視的習慣 一 被動吸煙 管教/限制 牙齒保健		
行為							
身體:		└──	正常 不正常	計 上			
整體外 皮膚		胸腔肺部		計	評估/ 計畫:		
頭部		心血管/脈搏		_			
耳朵	夏蓋物/ 無覆蓋物	腹部 生殖器		_			
鼻子 口咽/刃	- 齒	□					
頸部 淋巴結		神經步態					
請説明異常發現,並請加以評論。							
				已完	成的免疫接種		
			轉介	轉介			
下次檢查:3歳			保健	保健服務機構名稱			
保健服務機構簽名			保健	保健服務機構地址			
DSHS 13-684A CH (REV. 08/2004)							

童年早期:2歲

2歲小兒的保健

重要標誌

小兒在2至3歲期間的生長發育。

原地蹦跳。

騎三輪車。

説出帶三、四個字的句子。

自己吃飯和穿衣。

會用蠟筆劃十字和圓圈。

與娃娃和絨布動物玩"過家家"的游戲。

您可以在與您的孩子玩耍時幫助她學會新 技能。

若需要幫助或詳細資訊,請電

槍枝安全保存熱線:

1-800-LOK-IT-UP (565-4887)

兒童保健:

華盛頓州兒童保健服務資源與轉介網絡 1-800-446-1114

預防中毒:

華盛頓州防毒中心,1-800-732-6985或1-800-572-0638(TTY中繼轉發)

若您對您的子女感到很灰心,並需要幫助:

家庭幫助專線

1-800-932-HOPE (4673)當地危機熱線

子女教養技巧或支持:

家庭幫助專線 1-800-932-HOPE (4673)

西北地區家庭服務資源

1-888-746-9568當地社區學院課程

保健須知

您孩子最近是否定期接受過免疫接種?若是,他在 進幼稚園前就不需要再接受免疫接種。

每天為孩子提供各種不同的健康食品。少吃劣質食品。全家盡量經常在一起吃飯,並且吃飯時要關掉電視。

每天應該使用含氟牙膏至少為您的孩子刷一次牙, 牙膏用量應為豆粒大小。一定要為她每年做一次牙 科檢查。

子女教養須知

與孩子談論他所做的事。一起讀圖畫書並談論圖畫 的內容。

幫助孩子玩有一定運動量的游戲,如捉人游戲、捉 迷藏。給她簡單的玩具玩耍,如積木、蠟筆和紙 張、絨毛動物。

您可能想儘快訓練孩子自己大小便,但他可能要等到3歲時才可以做到。若他睡醒覺後尿布依然乾爽 並告訴您他想坐大小便,這就説明他可以開始學自己大小便了。

把兒童的電視時間限制到一小時或更短。與他們一 起觀看並談論節目的內容。

安全須知

- 將清潔用品鎖好並放在您的孩子接觸不到的 地方。
- 在靠近有交通往來的地方步行時(包括停車場),應始終拉著您孩子的手。在倒車前應先察看車後的情況,以防有兒童在車後方。
- 若您家中存有槍枝,應始終保存在安全處,並 要保證槍枝無子彈上膛並且鎖好。
- 每當孩子在水邊或船上時,均應穿上救生衣。 應始終在附近守望她。
- 把火柴和打火機放在您的孩子接觸不到的 地方。